

MISCELLANEOUS PAPERS

Centre for Entomological Studies Ankara

No 189

25 April 2019

ISSN 1015-8235

<http://zoobank.org/References/69BAB2EF-4CD1-4C50-9AE1-0F48996A834E>

Comments on the taxonomic status of “*marchandi* De Freina” in SE Turkey (Lepidoptera, Arctiidae)

Sibel Kızıldağ¹ Muhabbet Kemal Ahmet Ömer Koçak

Abstract: Comments on the taxonomic status of “*marchandi* De Freina” in SE Turkey (Lepidoptera, Arctiidae). *Misc. Pap.* 189: 1-3, 1 fig 2 Tables.

This paper deals with the molecular evaluation of the taxa *Arctia villica* and *marchandi* in SE Turkey. As a result of this, the name *marchandi* is proposed as a valid subspecific name of the *Arctia villica* populations of SE Turkey.

Key words: *Arctia*, *villica*, *marchandi*, *festiva*, Arctiidae, Lepidoptera, Turkey, mtCOI, genetic distance.

The genus *Arctia* was established by Schrank in 1802 (Fauna boica 2 (2): 152). Its type-species *Phalaena caja* Linnaeus, 1758 was designated by Westwood in 1840, subsequently. The name *marchandi*, proposed by De Freina (1983) with the name combination “*Arctia villica* ssp. *marchandi*” from “Zab-Tal” Zab Valley (Hakkari Province). *Marchandi* was used as valid name in various papers as a subspecies of *Arctia villica* (Linnaeus, 1758). De Freina (2011) published a revisional work on *Arctia villica*-complex. He proposed there *marchandi* as a distinct species, by basing upon the genital morphologic and genetic differences to *A. villica*.

In this paper, the mtCOI barcodes (658bp) of *Arctia villica*, *Arctia marchandi* and *Arctia festiva* populations from various localities in South and East Turkey have been evaluated. Among these taxa *Arctia villica* and *Arctia festiva* are well established, distinct species. The last name *marchandi* was suggested by De Freina (2011) as a distinct species. So far, the value of the genetic distance between *villica* and *marchandi* given by De Freina has not been confirmed in a publication. In fact, there is no clear, numerical calculation on the genetic distance between *villica* and *marchandi* in De Freina’s (2011) paper. It is almost impossible to understand and calculate the distance properly, by looking to Freina’s (2011: 97) tree diagram. This situation inevitably leads to some doubts. Consequently, it resulted in a discussion of the molecular evaluation and phylogenetic position of *Arctia villica*.

The mtCOI gene sequences in BOLDsystems and CESA BarcodingBank (**Table 2**) were selected for phylogenetic estimation of *Arctia villica*. Genetic distances between the populations were calculated based on the Kimura 2-parameter (K2P) model and phylogenetic relationships were assessed using the neighbour-joining (NJ) algorithm to construct gene trees with 1,000 replicates to evaluate internal nodes of the trees in MEGA6.

¹ Dr. Sibel Kızıldağ & Dr. Muhabbet Kemal - Van Yüzüncü Yıl University, Faculty of Science, Dept. of Biology, Campus, Van / Turkey.
e-mails: sibelkizildag@yyu.edu.tr - muhabbet_kemal@yahoo.com.tr
Prof. em. Dr. Ahmet Ömer Koçak, c/o Yüzüncü Yıl University, Faculty of Science, Dept. of Biology, Van / Turkey.
e-mail: cesa_tr@yahoo.com.tr




Fig. 1 - Kimura 2-parameter (K2P) distance neighbour-joining tree of COI gene sequences of *A.villica*, *A.villica marchandi* and *A. festiva* (outgroup) with bootstrap values.

In our phylogenetic tree populations of *Arctia villica* and *Arctia marchandi* were clustered in Clade A and Clade B (**Fig. 1**). Clade A was consist of *Arctia villica* and *Arctia marchandi* populations, Clade B was only consist of *Arctia marchandi* populations from East Turkey. Hakkari populations of *Arctia marchandi* have a sister position with *Arctia villica* populations in Clade A instead of populations in Clade B. This clearly shows that *Arctia villica/marchandi* is not a monophyletic group. Genetic distances based on K2P model were showed in **Table 1**. The K2P model genetic distances between populations of *Arctia villica* and *Arctia marchandi* is ranging from 0.2% to 1.4%. In Lepidopteran an empirical 2% (K2P) intraspecific distance value has often been proposed. In the present study, DNA barcode divergence between controversial pair of species was lower than 1.4%.

Table 1 - Percent pairwise corrected (K2P) genetic distance among the *Arctia villica*, *A. marchandi* and *A. festiva* COI sequences

1.	Arctia marchandi Hakkari Sumbul Arct 01																			
2.	VNMB690-08 Arctia marchandi COI-5P HQ684219	0.002																		
3.	CGUKD205-09 Arctia villica COI-5P KX043736	0.009	0.011																	
4.	Arctia marchandi Van Bahcesaray Arct 18	0.011	0.012	0.005																
5.	Arctia marchandi Mutki Aydemir Arct 15	0.011	0.012	0.005	0.000															
6.	Arctia marchandi Mutki Caygecit Arct 16	0.011	0.012	0.005	0.000	0.000														
7.	Arctia marchandi Mutki Kavakbasi Arct 17	0.011	0.012	0.005	0.000	0.000	0.000													
8.	GWORZ116-10 Arctia villica COI-5P HM914022	0.011	0.012	0.002	0.006	0.006	0.006	0.006												
9.	VNMB685-08 Arctia villica COI-5P HQ684224	0.011	0.012	0.011	0.012	0.012	0.012	0.012	0.012											
10.	Arctia marchandi Van Erek Arct 14	0.011	0.012	0.005	0.000	0.000	0.000	0.000	0.006	0.012										
11.	Arctia marchandi Van Agarti Arct 02	0.012	0.014	0.006	0.002	0.002	0.002	0.002	0.008	0.014	0.002									
12.	Arctia villica Anamur Arct 04	0.012	0.014	0.009	0.011	0.011	0.011	0.011	0.011	0.011	0.012									
13.	FBLMZ558-12 Arctia villica COI-5P	0.012	0.014	0.012	0.014	0.014	0.014	0.014	0.014	0.011	0.014	0.016	0.006							
14.	VNMB686-08 Arctia villica COI-5P HQ684223	0.014	0.016	0.008	0.012	0.012	0.012	0.012	0.009	0.016	0.012	0.014	0.014	0.017						
15.	Arctia festiva Mus Varto Arct 10	0.065	0.065	0.061	0.060	0.060	0.060	0.060	0.063	0.063	0.060	0.061	0.070	0.067	0.069					
16.	Arctia festiva Van Bahcesaray Arct 09	0.070	0.070	0.067	0.065	0.065	0.065	0.065	0.069	0.069	0.065	0.067	0.072	0.069	0.074	0.019				

Genetic distances between populations were not clearly reported in Freina's paper. Taking the K2P model genetic distances between populations of *Arctia villica* and *Arctia marchandi*, ranging from 0.2% to 1.4% into consideration, we propose here *marchandi* as a valid subspecific name of *Arctia villica*, as in the original form (**ssp. rev.**)

Table 2 – List of collecting places and their process ID of the the taxa used in tree diagram

Name of taxa under examination	Collecting data from Turkey	Process ID (Fasta)
<i>Arctia villica ssp. marchandi</i>	TR – Hakkari Prov., Sümbül Mt 1350m, M.Kemal & A.Koçak leg.	Lep-Arct01, 658bp
<i>Arctia villica ssp. marchandi</i>	TR – Van Prov., Ağartı, M.Kemal & A.Koçak leg.	Lep-Arct02, 658bp
<i>Arctia villica villica</i>	TR – İçel Prov., Anamur, M.Kemal & A.Koçak leg.	Lep-Arct04, 658bp
<i>Arctia festiva</i>	TR- Van Prov.,, Bahçesaray, Vasting, M.Kemal & A.Koçak leg.	Lep- Arct09, 658bp
<i>Arctia festiva</i>	TR- Muş Prov., Varto, Armutkaşı, M.Kemal & A.Koçak leg.	Lep- Arct10, 658bp
<i>Arctia villica ssp. marchandi</i>	TR- Van Prov., Ereğ Mt., M.Kemal & A.Koçak leg.	Lep- Arct14, 658bp
<i>Arctia villica ssp. marchandi</i>	TR- Bitlis Prov., Mutki, Aydemir, M.Kemal & A.Koçak leg.	Lep- Arct15, 658bp
<i>Arctia villica ssp. marchandi</i>	TR- Bitlis Prov., Mutki, Çaygeçit, M.Kemal & A.Koçak leg.	Lep- Arct16, 658bp
<i>Arctia villica ssp. marchandi</i>	TR- Bitlis Prov., Mutki, Kavakbaşı, M.Kemal & A.Koçak leg.	Lep- Arct17, 658bp
<i>Arctia villica ssp. marchandi</i>	TR- Van Prov., Bahçesaray, Vasting, M.Kemal & A.Koçak leg.	Lep- Arct18, 658bp

References

- Felsenstein J.**, 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39: 783-791.
- Freina, J.J. de**, 1983. 4. Beitrag zur systematischen Erfassung der Bombyces- und Sphinges-Fauna Kleinasiens. Neue Kenntnisse über Artenspektrum, Systematik und Nomenklatur sowie Beschreibung neuer Taxa (Lepidoptera). *Mitt. münch. Ent. Ges.* 72: 57–127.
- Freina, J.J. de**, 2011. Revision des *Arctia villica*-Komplexes 2. Teil: Nomenklatur und Systematik der osteuropäisch-asiatischen Arten *Arctia villica* (Linnaeus, 1758) (partim), *Arctia marchandi* De Freina, 1983 stat. n. und *Arctia confluens* Romanoff, 1884 stat. n. (Lepidoptera: Arctiidae, Arctiinae). *Entomofauna* 32 (4): 93-122.
- Kimura M.**, 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16:111-120.
- Nye, I.W.B.**, 1975. *The generic names of moths of the World*. Vol. I Noctuoidea (part). Trustees of the British Museum (Natural History).
- Saitou N. & M. Nei**, 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406-425.
- Tamura K., Stecher G., Peterson D., Filipowski A., & S. Kumar**, 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30: 2725-2729.

MISCELLANEOUS PAPERS**ISSN 1015-8235**

Miscellaneous Papers is a peer reviewed serial of the Centre for Entomological Studies Ankara, established in 1989. It appears at irregular interval in a year, as PDF format and includes original articles of the research workers of the Centre, regarding on various subjects on Entomology (taxonomy, nomenclature, checklist, fauna, biodiversity, distribution, biogeography, ecology, insect-plant interactions, bionomy, and behaviour).

The CESA is a non-profit group, no royalties will be paid to authors of contributions. Papers accepted become the copyright of the related serial.

Miscellaneous Papers is currently archived online at “Internet Archive”, in accordance with the publication rules of the ICZN. It is an open-access serial, distributed under the terms of the “Creative Commons Attribution License”, which permits free use, and distribution in any medium, provided the original author(s) and source are credited.

Centre for Entomological Studies Ankara



(A scientific Consortium)

(co-operation of research workers for pure-scientific, not commercial purpose)

Web Page of the Cesa: <http://www.cesa-tr.org/>

Scientific Serials: Priamus & Priamus Supplement (print and online versions) (ISSN 1015-8243)², Miscellaneous Papers (print and online versions) (ISSN 1015-8235)³, Memoirs (print and online versions) (ISSN-8227)⁴, DVD Films⁵, Iconographia Insectorum⁶ (online), Cesa Publications on African Lepidoptera (online)⁷, Cesa News (online)⁸, Cesa Books (online)⁹

Owners / Sahipleri - Editors / Yayıncılar: Prof. em Dr Ahmet Ömer Koçak (c/o Van Yüzüncü Yıl University, Van, Turkey), Asst. Prof. Dr Muhabbet Kemal Koçak (c/o Van Yüzüncü Yıl University, Van, Turkey).

Editorial Board of all Scientific Serials of the CESA / Bütün Bilimsel Yayınların Yayın Kurulu: Insecta, taxonomy, nomenclature, ecology, faunistics: Prof. em Dr Ahmet Ömer Koçak (c/o Van Yüzüncü Yıl Üniversitesi, Turkey), Asst. Prof. Dr Muhabbet Kemal Koçak (Van Yüzüncü Yıl University, Turkey).

Chief referees of all Scientific Serials of the CESA: Prof. em Dr Ahmet Ömer Koçak & Asst. Prof. Dr Muhabbet Kemal Koçak: Insecta, taxonomy, nomenclature, fauna, ecology, catalogues, checklists of the Old World.

Expert referees according to the subject areas: Dr Peter Huemer (Austria) and Dr. Ole Karsholt (Denmark): Gelechioidea, and some other Microlepidoptera groups in Palaearctic (Lepidoptera). Mag. Peter Buchner (Austria): Depressariidae (Lepidoptera). Dr J. B. Heppner (U.S.A.): Microlepidoptera of Nearctic and Neotropical. Dr G. Balduzzi (Italy): Coleophoridae (Lepidoptera). Dr V. Korneyev (Ukraine): Tephritidae, Pyrgotidae, Ulidiidae (Diptera). Prof. Dr Y.G.Verves (Ukraine): Sarcophagidae (Diptera). Dr Daniel Burckhardt (Switzerland): Psyllidae (Homoptera). Prof. Dr E. Heiss (Austria): Hemiptera. Dr R. Ehrmann (Germany): Mantodea. Prof. Dr Mustafa Ünal (Bolu, Turkey): Orthoptera. Prof. Dr Hüseyin Özdikmen (Turkey): Coleoptera. Prof. Dr Suat Kiyak (Turkey): Hemiptera.

Plant taxonomy, flora and vegetation: Assoc. Prof. Dr Murat Ünal, Asst. Prof. Dr Mesut Pınar, (Yüzüncü Yıl University, Van, Turkey).

Molecular studies: Asst. Prof. Dr Sibel Kızıldağ (Van Yüzüncü Yıl University, Van, Turkey).

Editorial policy: the submitted manuscript is evaluated by the Chief Editor and Referee. In case of need, the manuscript is sent to expert referees according to the subject areas.

ALL RIGHTS RESERVED

Correspondences should be addressed to: Prof. em Dr Ahmet Ömer Koçak, c/o Yüzüncü Yıl University, Fen Fakültesi, Biyoloji Bölümü, Kampus, Van / Turkey. - e-mail: cesa_tr@yahoo.com.tr

All the serials of the Cesa are archived regularly by the Internet Archive (300 Funston Ave., San Francisco, CA 94118, U.S.A.), in accordance with the rules of the International Codes of Zoological Nomenclature (ICZN)

<https://archive.org/>

also recorded regularly by the Zoological Record, Thomson Reuters, Enterprise House, Innovation Way, Heslington, York, YO10 5NQ, United Kingdom, <https://clarivate.com/>

² <http://www.cesa-tr.org/Pri.htm>

³ <http://www.cesa-tr.org/Miscell.htm>

⁴ <http://www.cesa-tr.org/Memoirs.htm>

⁵ <http://www.cesa-tr.org/CDF.htm>

⁶ <http://www.cesa-tr.org/Icon.htm>

⁷ <http://www.cesa-tr.org/CPAL.htm>

⁸ <http://www.cesa-tr.org/Cesanews.htm>

⁹ <http://www.cesa-tr.org/Cesabooks.htm>